arXiv:cond-mat/9803011v1 [cond-mat.supr-con] 2 Mar 1998

ON TEMPERATURE *versus* DOPING PHASE DIAGRAM OF HIGH CRITICAL TEMPERATURE SUPERCONDUCTORS

V.M. Loktev

Bogolyubov Institute for Theoretical Physics,

Metrologichna str. 14-b, Kyiv, 252143 Ukraine

V.M. Turkowski

Shevchenko Kyiv University

Acad. Glushkova prosp. 6, Kyiv, 252127 Ukraine

Abstract

The attempt to describe the bell-shape dependence of the critical temperature of high-T_c superconductors on charge carriers density is made. Its linear increase in the region of small densities (underdoped regime) is proposed to explain by the role of the order parameter phase 2D fluctuations which become less at this density growth. The critical temperature suppression in the region of large carrier densities (overdoped regime) is connected with the appearance (because of doping) of the essential damping of long-wave bosons which in the frame of the model proposed define the mechanism of indirect inter-fermion attraction.

PACS: 67.20.+k, 74.20.-z, 74.25.-q, 74.72.-h

1 Introduction

In spite of non-falling down the intensity of study of high-T_c superconductors (HTSCs) the comprehension of many regularities which define their physical behaviour is not still achieved. First of all, it is put down to the HTSC normal state properties, the description of which in the framework of standard Fermi-liquid theory proves to be impossible if the carrier concentration n_f in corresponding samples is like that one that the resulting critical temperature T_c in them is less than some characteristic for every compound optimal value T_c^{max}. The latter in all (practically, without exclusion) HTSC copper oxides appears because, as it is well-known (see, for example, the rewiev [1]), the conductivity (consequently - superconductivity) these, initially quasi-2D antiferromagnetic insulators, results from their doping by donor (Nd, Pr) or acceptor (Sr, Ba, O) ions.

Then the temperature T_c being formed at increasing of carrier density (or, what is the same, Fermi-energy) in a system also grows up. This growth, however, is rather quickly stopped and after some (not large upon n_f values) part of "saturation" the drop to zero of the function $T_c(n_f)$ follows, which thus acquires the bell-shape form. As a result, the HTSC compounds with relatively large concentration of itinerant carriers (so called, overdoped regime) become non-superconducting metals the behaviour of which to a certain degree can be carried out on the base of conventional theory of Fermi-liquid [2]. The corresponding carrier concentration values for underdoped regime are such ones that the $T_c(n_f)$ increase is observed for n_f from $\simeq 0.04 - 0.08$ carrier per CuO_2 layer cell (i.e. from the threshold of the insulator-metal transition) to $T_c^{max} \simeq T_c(0.15 - 0.18)$; the temperature $T_c(n_f)$ becomes zero when $n_f \simeq 0.25 - 0.28$ [3, 4, 5]. If the initial part of the function $T_c(n_f)$, where $dT_c(n_f)/dn_f \simeq const > 0$ [6, 7], can be (at least qualitatively) interpreted (see Refs.[8, 9, 10] and the review [11]) proceeding from the theory of the crossover from the Bose-Einstein condensation of separate composite bosons (local pairs) and their superfluidity to the superconductivity of the BCS type, then the reason of $T_c(n_f)$ suppression in the BCS carrier concentration region remains, in fact, unknown yet.

From the very begining the above-mentioned behaviour of $T_c(n_f)$ was ascribed to the strong electron-electron correlations (see, for example, the book [12]) and to the filling of usually narrow Hubbard subbands by itinerant carriers. This question was intensively investigated by Prof. I.V. Stasyuk and his collaborators [13, 14, 15, 16].

However a little bit later it becomes evident that carrier concentrations corresponding to a disappearance of superconductivity are so small that such a filling of the whole valent (in fact, conduction) band in HTSCs is impossible. The alternative version of the HTSC effect disappearance most consistently considered in Refs.[17, 18] (see also [1, 2]) consists in assumption that bare Fermi-level in HTSCs proves to be in the vicinity of the extended saddle point of the electronic spectrum. According with this poin of view at n_f growth the Fermi-energy $\epsilon_F \simeq \mathbf{k}_F^2/2m$ ($\mathbf{k}_F$ - Fermi momentum, m - effective mass) passes the van Hove singularity in the conduction band density of states and, hence, $T_c(n_f)$ reveals its maximum. If it would be so the function $T_c(n_f)$, being similar to BCS one, would have the exponential dependence on n_f far from T_c^{max} what, as it is seen from experiments and was already said, is not confirmed by observations. On the other hand, the very appearance of the van Hove spectrum singularity requires fine-tuning (for example, because of rather definite ratio between the hole next-near and nearest hopping constants) what seems can be hardly achieved even in one compound not speaking of many ones.

At the same time less attention was paid to the idea that the weakness of superconductivity can be conditioned by "feedback" effect, or the changes in the spectrum $\omega(\mathbf{k})$ of intermediate bosons the exchange of which results in the fermion pairing. These bosons in HTSCs can be attributed to phonons (similarly to BCS-Eliashberg model) the role of which is actively advocated by Ginzburg (see, for example, his rewiev [19]), magnons, or spin fluctuations, [20, 21], quadrupole dd-excitons (Gaididei-Loktev-Weber mechanism (see Ref.[1])), plasmons [22], etc. It must be, however, noted that if phonon spectrum does not (or very weakly) depend on doping then long-wave excitations of magnetic subsystem (dd-excitons, as excitations over a magnetic background, can be also considered here) are strongly suppressed because of the long-range magnetic order destruction in metallic phase of HTSCs [1, 12].

Namely spin waves are subjected the most appreciable doping effect; their low-frequency region (for the wave vectors less than some characteristic value k_{min}) acquires diffusion form, or becomes overdamped. In the insulating phase of HTSCs as it is shown in Ref. [23], $k_{min} \sim n_f$; in their mettalic phase the magnon damping becomes even more strong and $k_{min} \simeq 2k_F \sim n_f^{1/2}$ [24, 1], what is completely in line with degradation of the correlation magnetic length $\xi_{mag} \sim n_f^{-1/2}$ measured in HTSCs [25]. In that way the long-wave damping $\gamma(\mathbf{k})$ increases up to so high values ($\gamma(\mathbf{k}) >> \omega(\mathbf{k})$) that corresponding intermediate bosons (most probably, spin fluctuations) can not participate in interaction transfer, "being out".

The solution of the self-consistent magneto-electronic problem as a whole (i.e. the explicit solution of the equation of superconductivity similar to the Eliashberg equation) is scarcely possible now. Therefore below the attempt is made to consider the simplest model with indirect iner-fermion attraction what is provided by intermediate massive bosons with long-wave cut which is proportional to k_F. The carriers to be supposed appear in a system due to doping what corresponds to generally accepted scenario of metallization of copper oxides. For the sake of simplicity we shall omit any other dampings (in particular, the carrier damping because of disorder the effect of which was analysed, for example, in Ref.[26]).

Following to Refs.[9, 11] we shall also suppose that in a model 2D system (in fact, all HTSCs can be with a good accuracy put down to this kind) the superconducting condensate is formed in a way principally different from the ordinary one. In such a case one has to differ the order parameter formation temperature from the real (observable) critical temperature in a sense that the absolute value of the former does not become zero at and above T_c.

2 Model and main equations

The model Hamiltonian density of the electron-phonon system can be written in the well-known form:

$$\begin{aligned} H(x) = -\psi_\sigma^\dagger(x)\left(\frac{\nabla^2}{2m} + \mu\right)\psi_\sigma(x) + H_{ph}(\varphi(x)) \\ + g_{ph}\psi_\sigma^\dagger(x)\psi_\sigma(x)\varphi(x), \quad x = \mathbf{r}, t, \end{aligned} \tag{1}$$

where H_{ph} is the Hamiltonian of free phonons which will be described more precisely below; $\psi_\sigma(x)$ and $\varphi(x)$ are the fermion and boson field operators, respectively; m is the effective mass of the fermi-particles, $\sigma =\uparrow, \downarrow$ - their spin variable, and g_{ph} is the electron-phonon coupling constant; μ in (1) is the chemical potential of fermions, which fixes their average density in a system; we put $\hbar = k_B = 1$.

As it was mentioned in Introduction, we shall model boson-exchange interaction taking into account the dependent on doping saturation of long-wave bosons. Under above mentioned assumptions about gradual non-participation of the part of the bosons in an attracting interaction formation the simplest way to describe that can be achieved by making use of the free-phonon propagator in the form:

$$D(\omega, \mathbf{k}) = -\frac{\omega^2(\mathbf{k})}{\omega^2 - \omega^2(\mathbf{k}) + i\delta}\theta(k - k_{min}), \quad \delta \to 0, \tag{2}$$

where $\omega(\mathbf{k})$ is the boson dispersion law, and $\theta(k)$ is the step function. As it was pointed out above $\mathbf{k}_{min}$ in (2) is some characteristic wave vector which devides the region of overdamped ($k < k_{min}$) and long-lived ($k > k_{min}$) intermediate bosons (here: phonons by definition). Although magnetic correlation length measurements result in $k_{min} = 2k_F$ we shall adopt a more general (or soft) relation supposing that $k_{min} = \alpha k_F \equiv \alpha\sqrt{2m\epsilon_F}$ where α is some free-parameter.

It is very important that the Hamiltonian (1) is invariant with respect to the global symmetry transformations

$$\psi_\sigma(x) \to \psi_\sigma(x)e^{i\theta}, \quad \psi^\dagger_\sigma(x) \to \psi^\dagger_\sigma(x)e^{-i\theta}, \tag{3}$$

which in 2D case (unlike to 3D one) remain unbroken and phase transition is here accompanied by change in the correlation function behaviour only.

The $T{-}n_f$ phase diagram of a system can be calculated by using the Hubbard-Stratonovich method, generalized on the case of non-local (indirect) interaction (so called auxiliary bilocal field method). For the finding of the grand partition function Z it is useful to pass to Nambu spinors: $\Psi^\dagger = (\psi^\dagger_\uparrow, \psi_\downarrow)$ and its conjugated one. After that performing the integration over bosonic fields it is easy to obtain the Lagrangian

$$\begin{aligned} L = \Psi^\dagger(x)[-\partial_\tau + (\frac{\nabla^2}{2m} + \mu)\tau_z]\Psi(x) \\ -\frac{1}{2}\Psi(x_1)\Psi^\dagger(y_1)\tau_z K(x_1, y_1; x_2, y_2)\Psi(x_2)\Psi^\dagger(y_2)\tau_z \end{aligned} \tag{4}$$

of a system, where the integration over repeated indices is supposed. The kernel K in (4) will be defined below.

Let introduce the pairing order parameter

$$\phi(x_1,y_1) = K(x_1,y_1;x_2,y_2)\Psi(x_2)\Psi^\dagger(y_2)\tau_z$$

$$\equiv \tau_+\phi(x_1,y_1) + \tau_-\phi^*(x_1,y_1), \tag{5}$$

where $\tau_+ = \frac{1}{2}(\tau_x + i\tau_y)$, $\tau_- = \frac{1}{2}(\tau_x - i\tau_y)$ (and τ_z in (4)) are the Pauli matrices.

Then adding to the Lagrangian L a zero term

$$\frac{1}{2}[\phi(x_1,y_1) - K(x_1,y_1;x_1',y_1')\Psi(x_1')\Psi^\dagger(y_1')\tau_z]K^{-1}(x_1,y_1;x_2,y_2)[\phi(x_2,y_2)$$

$$-K(x_2,y_2;x_2',y_2')\Psi(x_2')\Psi^\dagger(y_2')\tau_z]$$

with the purpose to cancel the four-fermion interaction one comes to the expression:

$$L(x_1,y_1;x_2,y_2) = \Psi^\dagger(x_1)[-\partial_\tau + (\frac{\nabla^2}{2m} + \mu)\tau_z - \tau_+\phi(x_1,y_1)$$

$$-\tau_-\phi^*(x_1,y_1)]\Psi(y_1) + \frac{1}{2}\phi(x_1,y_1)K^{-1}(x_1,y_1;x_2,y_2)\phi(x_2,y_2) \tag{6}$$

for the Lagrangian needed. The Fourier transformation of K can be written as

$$K(x_1,y_1;x_2,y_2) = \int \frac{d^3Pd^3p_1d^3p_2}{(2\pi)^9}K_P(p_1;p_2)exp[-iP(\frac{x_1+y_1}{2} - \frac{x_2+y_2}{2})$$

$$-ip_1(x_1-y_1) - ip_2(x_2-y_2)],$$

($p_i = (\mathbf{p}_i, \omega_i)$ where $i = 1,2$ and $P = (\mathbf{P}, \omega)$ are the relative and the centre of mass momenta, respectively). Supposing now that $K_P(p_1;p_2)$ is P- independent we pass to the standard kernel form

$$K(p_1;p_2) = g_{ph}^2 D(p_1 - p_2), \tag{7}$$

which corresponds to the indirect inter-fermion interaction.

The partition function can be written then as

$$Z = \int \mathcal{D}\Psi^\dagger\mathcal{D}\Psi\mathcal{D}\phi\mathcal{D}\phi^* \exp\left[-\beta\int L(\Psi^\dagger,\Psi,\phi^*,\phi)dxdy\right]$$

$$\equiv \int \mathcal{D}\phi\mathcal{D}\phi^* \exp(-\beta\Omega[\mathcal{G}]), \quad \beta \equiv 1/T,$$

In the last expression $\Omega[\mathcal{G}]$ is the thermodynamic potential which in the "leading order" on g_{ph} has the form

$$\beta\Omega[\mathcal{G}] = -\mathrm{Tr}\left[\mathrm{Ln}\mathcal{G}^{-1} + \frac{1}{2}(\phi K^{-1}\phi)\right], \tag{8}$$

in which

$$\mathcal{G}^{-1} = -\left[\partial_\tau - \left(\frac{\nabla^2}{2m} + \mu\right)\tau_z - \tau_+\phi - \tau_-\phi^*\right] \tag{9}$$

is the full fermion Green function. After the direct minimization of the potential (8) it is easy to obtain the equation for auxiliary ϕ-field (or order parameter):

$$\delta\Omega/\delta\phi = 2\phi - tr\int\frac{d^2\mathbf{k}d\omega}{(2\pi)^3}K(p;\mathbf{k},\omega)\mathcal{G}(\mathbf{k},\omega)\tau_x = 0. \tag{10}$$

Using (10) one can readily arrive to the well-known Cornwell-Jackiw-Tomboulis formula for the effective action in the one-loop approximation [27]:

$$\beta\Omega(\mathcal{G}) = -\mathrm{Tr}\mathrm{Ln}\mathcal{G}^{-1} + \frac{1}{2}\mathrm{Tr}\mathcal{G}K\mathcal{G},$$

or, taking into account (9) (or (10)),

$$\beta\Omega(\mathcal{G}) = -\mathrm{Tr}\left[\mathrm{Ln}\mathcal{G} + \frac{1}{2}(\mathcal{G}\mathcal{G}_0^{-1} - 1)\right]. \tag{11}$$

To investigate the possibility of the condensate formation in a 2D system it is convenient in accordance with [9] to pass to a modulus-phase parametrization of the order parameter (cp. (3), where $\theta = const$):

$$\phi(x,y) = \rho(x,y)exp[-i(\theta(x) + \theta(y))/2] \tag{12}$$

with simultaneous Nambu spinor transformation

$$\Psi^\dagger(x) = \chi^\dagger(x)exp[i\theta(x)\tau_z/2], \tag{13}$$

corersponding to "separation" of the bare fermions on their neutral $\chi(x)$ and charge $\theta(x)$ parts (fermi- and bose-ones, respectively).

In the approximation that $\rho(x,y) = \rho = const$ (see Ref.[28]) and spatial θ-fluctuations are small one can (using (12) and (13)) obtain the next expressions for $\mathcal{G}$ and Ω which are defined in (9) and (11):

$$\begin{aligned}\mathcal{G}^{-1} &= -\left[\partial_\tau - \tau_z\left(\frac{\nabla^2}{2m}+\mu\right)+i\tau_x\rho\right. \\ &\quad \left. -\tau_z\left(\partial_\tau\theta+\frac{\nabla\theta^2}{2m}\right)-i\left(\frac{\nabla^2\theta}{2m}+\frac{\nabla\theta\nabla}{m}\right)\right] \\ &\equiv G^{-1}(\rho)-\Sigma(\partial\theta) \qquad (14)\end{aligned}$$

and $\Omega = \Omega_{kin}(\rho, \nabla\theta) + \Omega_{pot}(\rho)$ with $\Omega_{pot}(\rho)$ which is defined by (11) at $\nabla\theta = 0$ and

$$\beta\Omega_{kin}(\rho,\nabla\theta) = \mathrm{Tr}[G\Sigma - G_0\Sigma + \frac{1}{2}G\Sigma G\Sigma - \frac{1}{2}G_0\Sigma G_0\Sigma$$

$$+ \tau_x\frac{1}{2}i\rho G(G\Sigma + G\Sigma G\Sigma)] = \frac{T}{2}\int_0^\beta d\tau\int d^2\mathbf{r}J(\mu,T,\rho(\mu,T))(\nabla\theta)^2, \qquad (15)$$

where the effective neutral fermion stiffness

$$\begin{aligned} &J(\mu,T,\rho(\mu,T)) \\ &= \frac{1}{8\pi}\left[\sqrt{\mu^2+\rho^2}+\mu+2T\ln\left(1+\exp(-\frac{\sqrt{\mu^2+\rho^2}}{T})\right)\right] \\ &- \frac{T}{4\pi}\left[1-\frac{\rho^2}{4T^2}\frac{\partial}{\partial(\rho^2/4T^2)}\right]\int_{-\mu/2T}^{\infty}dx\frac{x+\mu/2T}{\cosh^2\sqrt{x^2+\rho^2/4T^2}} \qquad (16)\end{aligned}$$

was introduced.

The evident analogy with XY-model (two-component order parameter in 2D space) gives the equation for the temperature T_{BKT} of the Berezinskii-Kosterlitz-Thouless phase transition in a system, namely (see Chapter 15 in the book [29]):

$$\frac{\pi}{2}J(\mu, T_{BKT}, \rho(\mu, T_{BKT})) = T_{BKT}. \qquad (17)$$

(Recall that the temperature T_{BKT} plays the role of critical one in 2D metals.)

The parameters μ and ρ in (17) being dependent of T are still unknown; therefore it is necessary to obtain the equations which connect them with carriers density n_f. The first one follows from (10) for $\rho \neq 0$:

$$1 = T\sum_{m=-\infty}^{\infty}\int\frac{d^2\mathbf{k}}{(2\pi)^2}\frac{K(\omega_m)}{\omega_m^2+\xi^2(\mathbf{k})+\rho^2}, \qquad (18)$$

where $\omega_n = (2n+1)\pi T$ is the Matsubara fermion frequencies, $\xi(\mathbf{k}) = \mathbf{k}^2/2m - \mu$ and it was used the Einstein model for phonon dispersion: $\omega(\mathbf{k}) = \omega_0$. The dependence on parameter α in Eq.(20) is preserved through the kernel K (see (7) and (2)).

The second one is defined by the condition $V^{-1}\partial\Omega_{pot}(\rho)/\partial\mu = -n_f$ (V is the volume of a system) what results in the well-known number equation:

$$\sqrt{\mu^2+\rho^2} + \mu + 2T\ln\left[1+\exp\left(-\frac{\sqrt{\mu^2+\rho^2}}{T}\right)\right] = 2\epsilon_F, \tag{19}$$

where the equality $\epsilon_F = \pi n_f/m$ was used; it is correct for free 2D fermions with a quadratic dispersion law.

Thus, we have obtained a self-consistent set of the Eqs.(17)-(19) needed to investigate the phase diagram of a 2D metal with arbitrary carrier density. The latter is the parameter defined such a metal superconducting properties.

3 Phase diagram of a system

As it can be seen from the previous Section (see also Refs.[9, 11]) there exist two characteristic temperatures in a system: T_ρ, where formally the complete order parameter given by Eq.(5) arises but its phase is a random quantity, i.e. $<\phi(x,y)>= 0$, and another one, $T_{BKT} < T_\rho$, where the phase of the order parameter becomes ordered, so that $<\phi(x,y)>\neq 0$. It must be, however, stressed that the temperature T_ρ is not a real critical temperature; it only denotes the characteristic region where the modulus of the order parameter achieves its maximal growth at T decreasing. Unlike T_ρ the temperature T_{BKT} does correspond to phase transition when correlators $< exp[i\theta(\mathbf{r},\tau)]exp[i\theta(\mathbf{r}')] >$ as function of $|\mathbf{r}-\mathbf{r}'|$ change their behaviour [29].

Let us find the n_f-dependence of the temperatures T_ρ and T_{BKT}. "Effective" temperature T_ρ can be estimated from the set (16)-(19) in mean-field approximation by putting $\rho = 0$. Another temperature T_{BKT} follows from the Eqs. (17) and (19).

It is impossible to solve the equations obtained analytically, so we shall do that by numerical calculation. Nevertheless, some assymptotical expressions for these temperatures as functions of n_f can be found in analytical form:

i) at $\epsilon_F/\omega_0 \to 0$ one obtains $T_{BKT} = \epsilon_F/8$, and T_ρ satisfies the simple mean-field equation $T_\rho \ln(T_\rho/\epsilon_F) = \omega_0 \exp(-2/\lambda)$, where $\lambda = g_{ph}^2 m/2\pi$ is the dimensionless coupling constant.

ii) $\epsilon_F \to \epsilon_F^{cr}$; the critical point $\epsilon_F = \epsilon_F^{cr}$ (or $n_f = n_f^{cr}$, what is the same) at which $T_\rho = T_{BKT} = 0$ can be found from Eqs.(17)-(19). This unknown energy is the solution of the equation

$$1 = \frac{\lambda}{2} \ln \frac{(W - \epsilon_F)[(\alpha - 1)\epsilon_F + \omega_0]}{(\alpha - 1)\epsilon_F(W - \epsilon_F + \omega_0)} \to \frac{\lambda}{2} \ln \frac{(\alpha - 1)\epsilon_F + \omega_0}{(\alpha - 1)\epsilon_F}|_{W\to\infty}. \tag{20}$$

(W is the conduction bandwidth determined by evident condition: $W = \mathbf{k}_B^2/2m$, where $\mathbf{k}_B$ is the Brillouin wave vector). In other words, it follows from (20) that because of the long waves phonons begin to quit the interaction transfer, there exist (at $\alpha > 1$ only), the point where both T_ρ and T_{BKT} temperatures become zero what means that superconductivity (but not conductivity) is suppressed. Near this point this temperatures have the next behaviour ($W \to \infty$):

$$T_\rho = \omega_0 / \ln \frac{4\epsilon_F^{cr}[(\alpha - 1)\epsilon_F^{cr} + \omega_0]}{\lambda\omega_0(\epsilon_F^{cr} - \epsilon_F)};$$

$$T_{BKT} = \epsilon_F^{cr} \left[\frac{9}{144}(\alpha - 1)^4 \left(\frac{\lambda\omega_0(\epsilon_F^{cr} - \epsilon_F)}{\epsilon_F^{cr}[(\alpha - 1)\epsilon_F^{cr} + \omega_0]} \right)^2 \right]^{1/5}.$$

The results of numerical calculations of Eqs. (17)-(19) are shown on Fig.1.

It is seen that due to the long-wave phonon suppression (in fact, to be supposed strong) the functions $T_\rho(n_f)$ and $T_{BKT}(n_f)$ quickly on n_f variation acquire a bell-like shape. This "bell" proves to be non-symmetrical, and its heigt, width and to some extent form depend on $\alpha(> 1)$. In some sense such a result is surprising because, as it is generally accepted, not long-, but short-wave intermediate bosons play the main role in attraction which appears in BCS-Eliashberg model due to electron-phonon (or any other boson) interaction. The sensitivity of superconducting properties of a 2D metal to long-wave part of the intermediate boson spectrum is rather unusual and allows to hope the more accurate consideration also results in a similar effect.

4 Conclusion

The existence of two-different temperatures in underdoped HTSCs is now a well established fact. They, as it shown in many papers (see the rewiev [11]), are one of the consequences of two-dimensionality of their electronic and magnetic properties when the ”ordering” of the order parameter modulus and phase takes place at different temperatures. In pure 2D system the temperature T_{BKT}, as was mentioned, has to be considered as critical one, and in the region $T_{BKT} < T < T_\rho$ so called pseudogap, also normal, phase is formed in underdoped HTSCs. It is destroyed when $T > T_\rho$ or n_f becomes so large (overdoped regime) that chemical potential of fermions and Fermi energy are indistinctive ($\mu = \epsilon_F$). And the aim of this paper was to demonstrate that in the superconducting system with indirect attraction the role of long-wave bosons (for instance, phonons, spin fluctuations and so on) can be crucial in the region where standard Fermi-liquid theory becomes already applicable.

In spite of some qualitative similarity between experimental and obtained pictures the model considered is so simple (and even rough because the propagator (2) was, in fact, postulated while it must contain the boson damping only) that any quantitative its use is almost senseless. Therefore it must be generalized on the taking into account such HTSC features as: quasi-two-dimensionality (which results in real T_c), intermediate boson dispersion $\omega(\mathbf{k})$ (as it takes place for spin fluctuations) and damping $\gamma(\mathbf{k})$; pairing anisotropy, etc. These problems will be considered separately.

Acknolegements

We are pleased to devote this paper to well-known ukrainian theorist, Corresponding member of the National Academy of Sciences of Ukraine, Prof. I.V. Stasyuk whose numerical remarks and critics are always interesting, useful and helpful for us.

References

[1] Loktev V.M. Mechanisms of high-T_c superconductivity of copper oxides.// Low Temp. Phys., 1996, vol.22 , No 1, p. 1-32.

[2] Shen Z.-X., Dessau D.S. Electronic structure and photoemission studies of late transition - metal oxides - Mott insulators and high-temperature super-conductors.// Phys. Rep., 1995, vol.253, No 1-3, p.1-162.

[3] Keimer B., Belk N., Birgeneau R.J., Cassano A., Chen C.Y., Greven M., Kastner M.A., Shirane G. Magnetic excitations in pure, lightly doped and weakly metallic La_2CuO_4.// Phys. Rev. B, 1992, vol.46, No 21, p.14034-14053.

[4] Takagi H., Cava R.J., Marezio M., Batlogg B., Krajewski J.J., Peck W.F., Jr., Bordet P., Cox D.E. Disappearance of superconductivity in overdoped $La_{2-x}Sr_xCuO_4$ at structural phase boundary.// Phys. Rev. Lett., 1992, vol.68, No 25, p.3777-3780.

[5] Nagano T., Tamioka Y., Nakayama Y., Kishio K., Kitazawa K. Bulk superconductivity in both tetragonal and orthorhombic solid solutions of $(La_{1-x}Sr_x)_2CuO_{4-\delta}$ at structural phase boundary.// Phys. Rev. B 1993, vol.48, No 13, p.9689-9696.

[6] Uemura Y.J., Le L.P., Luke J.M., Sternlieb B,J., Wu W.D., Brewer J.H., Riesman T.M., Seaman C.L., Maple M.B., Ishikawa M., Hinks D.G., Jorgensen J.D., Saito G., Yamchi H. Basic similarities among cuprate, bismthate, organic, Chevrel- Phase and heavy-fermion superconductors shown by penetration- depth measurements.// Phys. Rev. Lett. 1991, vol. 66, No20, p.2665-2668.

[7] Yamada K., Wakimoto S., Shirane G., Lee C.H., Kastner M.A., Hosoya S., Greven M., Endoh Y., Birgeneau R.J. Direct observation of a magnetic gap in superconducting $La_{1.85}Sr_{0.15}CuO_4$ ($T_c = 37.3$K).// Rev. Lett., 1995, vol. 75, No 8, p.1626-1629.

[8] Emery V., Kivelson S.A. Importance of phase fluctuations in superconductors with small superfluid density.// Nature 1995, vol. 374, p.434-437.

[9] Gusynin V.P., Loktev V.M., Sharapov S.G. Phase diagram of 2D metal with variable number of carriers.// JETP Lett., 1997, vol. 65, No 2, p.182-188; On peculiarities of

superconducting state formation in 2D metallic systems.// Low Temp. Phys., 1997, vol.23 , No 8, p.

[10] Abrikosov A.A. Size of BCS ratio in strongly underdoped high-T_c cuprates.// Phys. Rev. B 1997, vol. 55, No 10, p.R 6149-R 6151.

[11] Loktev V.M., Sharapov S.G. Superconducting condensate formation in metallic systems with arbitrary carrier density.// Cond. Mat. Phys., 1997, No 11, p.131-178.

[12] Plakida N.M. High-Temperature Superconductors, Berlin, Springer Verlag, 1995.

[13] Plakida N.M., Yushankhai V.Yu, Stasyuk I.V. On the role of kinematic and exchange interactions in superconducting pairing of electrons in the Hubbard model.// Physica C, 1989, vol. 160, No 1, p.80-88.

[14] Stasyuk I.V., Shvaika A.M., Schachinger E. On the electron spectrum of the Hubbard model including the interaction with loca anharmonic vibrations.// Physica C, 1993, vol. 213, p.57-70.

[15] Stasyuk I.V., Shvaika A.M. Dielectric properties and electron spectrum of the Müller model in HTSC theory.// Acta Phys.Polonica A, 1993, vol. 84, No 2, p.293-313.

[16] Stasyuk I.V., Shvaika A.M., Schachinger E. Electron spectrum of the Hubbard model with coupling to pseudospin degrees of freedom.// Physica B, 1994, vol. 194-196, p.1965-1966.

[17] Friedel J. The high-T_c superconductors: a conservative view.// J.Phys.: Cond. Mat., 1989, vol. 1, No 42, p. 7757-7794.

[18] Markiewicz R.S. Van Hove singularities and high-T_c superconductivity: A review.// Int. J. Mod. Phys. B, 1991, vol.5, No 12, p. 2037-2071.

[19] Ginzburg V.L. Superconductivity and superfluidity (what is done and what is not done).// Usp. Fiz. Nauk, 1997, vol. 197, No 4, p.429-454 (in Russian).

[20] Kampf A.P. Magnetic correlations in high-temperature superconductivity.// Phys. Rep., 1994, vol. 249, No 4,5, p.219-351.

[21] Pines D. Understanding high temperature superconductivity: progress and prospects.// CNLS Newsletter, Los Alamos Nat. Lab., 1997, No 138.

[22] Pashitskii E.A. Low frequency charge excitations in cuprate metal-oxide compounds// Fiz. Nizkikh Temp., 1995, vol. 21, No 10, p.995-1019; No 11, p.1091-1137 (in Russian).

[23] Ivanov M.A., Loktev V.M., Pogorelov Yu.G. Localization of spin excitations and disruption of long-range order in weakly doped La_2CuO_4.//JETP, 1992, vol.74, No 2, p.317-325.

[24] Loktev V.M. About correlation length in magnetic subsystem of high-T_c superconductors.// Supercond.: Physics, Chemistry, Technics, 1991, vol. 4, No 12, p.2993-2296.

[25] Thurston T.R., Birgeneau R.J., Gabbe D.R., Jenssen H.P., Kastner M.A., Picone P.J., Preyer N.W., Axe J.D., Böni P., Shirane G., Sato M., Fukuda K., Shamoto S. Neutron scattering study of soft optical phonons in $La_{2-x}Sr_xCuO_{4-y}$.// Phys. Rev. B, 1989, vol. 39, No 7, p.4327-4333.

[26] Loktev V.M., Pogorelov Yu.G. Metallization and superconductivity in doped metal-oxide layered compounds.// Physica C, 1996, vol. 272, p.151-160.

[27] Cornwell J.M., Jackiw R., Tomboulis E. Effective action for composite operators // Phys. Rev. D, 1974, vol. 10, No 8, P.2428-2445.

[28] Loktev V.M., Sharapov S.G., Turkowski V.M. Crossover from supefluidity to superconductivity in 2D systems with indirect inter-carrier interaction.// J. of Phys. Studies, 1997, vol. 1, No 3, p.431–440.

[29] Izyumov Yu.A., Skryabin Yu.N. Statistical mechanics of magnetically ordered systems. Moscow, Nauka, 1987 (in Russian).



Figure 1: The characteristic patterns of the $T-n_f$ phase diagram of 2D metal with coupling constant $\lambda = 1$. Doted and solid lines define the temperatures T_ρ and T_{BKT}, respectively.